

The signal of $Z^\pm(4430)$ in nucleon-antinucleon scattering

Hong-Wei Ke¹ and Xiang Liu^{2*†}

¹*Department of Physics, Nankai University, Tianjin 300071, China*

²*Centro de Física Computacional, Departamento de Física,
Universidade de Coimbra, P-3004-516 Coimbra, Portugal*

(Dated: October 2, 2008)

We study the production of $Z^\pm(4430)$ at a nucleon-antinucleon scattering experiment. Considering the PANDA experiment to be an ideal platform to explore the production of the charmonium and charmonium-like states, we suggest the forthcoming PANDA experiment to pay attention to the production of $Z^\pm(4430)$.

PACS numbers: 13.75.Cs

I. INTRODUCTION

Recently the Belle Collaboration announced the observation of $Z^\pm(4430)$ in the invariant mass spectrum of $\psi'\pi^\pm$ of $B \rightarrow K\psi'\pi^\pm$. As an excellent candidate of the exotic state, $Z^\pm(4430)$ possesses $m = 4433 \pm 4(\text{stat}) \pm 1(\text{syst})$ MeV and $\Gamma = 44^{+17}_{-13}(\text{stat})^{+30}_{-11}(\text{syst})$ MeV [1]. Up to now, theorists have paid extensive attention to $Z^\pm(4430)$ [2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17]. $Z^\pm(4430)$ was explained as an S-wave threshold effect [2], a $D_1\bar{D}^*/D_1'\bar{D}^*$ molecular state [3, 4, 5, 6, 7, 14], a tetraquark state [8, 9, 10, 15], a cusp effect [11], a $\Lambda_c - \bar{\Sigma}_c^0$ bound state [12] and the radially excited state of D_s [17]. Besides exploring the structure of $Z^\pm(4430)$, the hidden and open charm decays were studied in Ref. [3, 16].

Up to now, $Z^\pm(4430)$ was only observed by the Belle experiment. Since searching $Z^\pm(4430)$ in other experiments will be helpful to establish $Z^\pm(4430)$ finally, the theoretical study of the production of $Z^\pm(4430)$ is an interesting topic. For example, recently the author of Ref. [15] studied the production of $Z^\pm(4430)$ by the photo-production experiment.

Exploring the charmonium and charmonium-like states above the open charm threshold is one of the most important physics aims of the forthcoming PANDA (AntiProton Annihilations at Darmstadt) experiment at FAIR [18]. Recently some groups performed the phenomenological studies of the productions of the charmonium and charmonium-like states around the PANDA experiment. Barnes and Li calculated the differential and total cross sections of the near-threshold associated charmonium production processes $p\bar{p} \rightarrow \pi\Psi$ with $\Psi = \eta_c, J/\psi, \psi', \chi_{c0}, \chi_{c1}$ using a simple initial-state light meson emission model [19]. In Ref. [20], the authors studied the production of $X(3872)$ at the PANDA experiment assuming $X(3872)$ to be the molecular state $D\bar{D}^*$ or the charmonium state $\chi_{c1}(2P)$. These instructive explorations not only further guide the PANDA experiment to find charmonium or charmonium-like state but also

stimulate more theorists to focus on this field.

The PANDA experiment could be an ideal platform to explore the production of $Z^\pm(4430)$. Thus, in this work, we first carry out a phenomenological study of $Z^+(4430)$ production at antiproton-proton scattering experiment. Meanwhile we also tentatively investigate $Z^-(4430)$ production by the antiproton-neutron annihilation process.

This work is organized as follows. After the introduction, we present the formula of the production of $Z^+(4430)$ by the $p\bar{p}$ interaction. Next, we formulate the expressions of $Z^-(4430)$ production by the $\bar{p}n$ annihilation in Sec. III. Section IV is for numerical results and our discussion.

II. THE PRODUCTION OF $Z^+(4430)$ BY THE SCATTERING OF PROTON AND ANTIPROTON

In 1987, Gaillard, Maiani and Petronzio for the first time applied the hadron-level pole diagram to calculate the production of charmonium from the $p\bar{p}$ states with fixed quantum number [21]. Later Lundborg, Barnes and Wiedner developed this method to do in-depth study of $p\bar{p} \rightarrow \pi^0\Psi$ ($\Psi = \eta_c, J/\psi, \psi', \chi_{c0}, \chi_{c1}$) [19, 22]. The hadron-level pole diagram is a suitable approach to study the charmonium or charmonium-like state in the low-energy $p\bar{p}$ scattering process.

As an excellent candidate for the exotic state, the component of $Z^+(4430)$ should be $c\bar{c}q_1\bar{q}_2$ for explanations in terms of the $D_1\bar{D}^*$ molecular state [4, 5] or the tetraquark state [8]. Thus the $Z^+(4430)$ production mechanism for the $p\bar{p}$ scattering process is similar to $p\bar{p} \rightarrow \pi^0\Psi$. In this work, we use the so called hadron-level pole diagram to depict the production of $Z^+(4430)$ at the proton and antiproton scattering process, which is depicted in Fig. 1. By exchanging neutron between proton and antiproton, the process of $p\bar{p} \rightarrow Z^+(4430)\pi^-$ occurs.

In this work, we consider three possible quantum numbers for $Z^+(4430)$, i.e. $J^P = 0^-, 1^-, 2^-$ [4]. The effective Lagrangians describing the interaction of $Z^+(4430)$ with

*Corresponding author

†Electronic address: liuxiang@teor.fis.uc.pt




FIG. 1: The diagram depicting the production of $Z^+(4430)$ in the proton-antiproton scattering process.

nucleons are written as

$$\mathcal{L}_{Z^+p\bar{n}} = \begin{cases} g_{Zpn}^{(0)} \bar{n} \gamma_5 p Z & \text{for } J^P = 0^- \\ ig_{Zpn}^{(1)} \bar{n} \gamma_\mu p Z^\mu & \text{for } J^P = 1^- \\ g_{Zpn}^{(2)} \gamma_5 \partial_\mu \bar{n} \partial_\nu p Z^{\mu\nu} & \text{for } J^P = 2^- \end{cases} \quad (1)$$

The value of the coupling constants $g_{Zpn}^{(0),(1),(2)}$ will be given in the section on the numerical results. The Lagrangian relevant to the coupling of π and the nucleons is well established [23]:

$$\begin{aligned} \mathcal{L}_{\bar{N}N\pi} &= -i\mathcal{G}\bar{N}\gamma_5\boldsymbol{\tau}\cdot\boldsymbol{\pi}N \\ &= -i\sqrt{2}\mathcal{G}\left[\bar{p}\gamma_5n\pi^+ + \bar{n}\gamma_5p\pi^- + \frac{(\bar{p}\gamma_5p - \bar{n}\gamma_5n)\pi^0}{\sqrt{2}}\right], \end{aligned} \quad (2)$$

where $\bar{N} = (\bar{p}, \bar{n})$. The coupling constant $\mathcal{G}$ is taken to be 13.5.

Using the above Lagrangian, we write out the amplitudes of the $p\bar{p} \rightarrow Z^+(4430)\pi^-$ process depicted in Fig. 1

$$\mathcal{M}^{J=0} = \sqrt{2}\mathcal{G}g_{pnZ}^{(0)}\bar{v}_{\bar{p}}\gamma_5\frac{1}{\not{p}_2 - \not{p}_b - m_n}\gamma_5u_p, \quad (3)$$

$$\mathcal{M}^{J=1} = \sqrt{2}i\mathcal{G}g_{pnZ}^{(1)}\bar{v}_{\bar{p}}\gamma_5\frac{1}{\not{p}_2 - \not{p}_b - m_n}\gamma_\mu u_p\varepsilon^\mu, \quad (4)$$

$$\mathcal{M}^{J=2} = -\sqrt{2}\mathcal{G}g_{pnZ}^{(2)}\bar{v}_{\bar{p}}\gamma_5\frac{1}{\not{p}_2 - \not{p}_b - m_n}\gamma_5u_p p_a^\mu p_b^\nu \varepsilon_{\mu\nu}, \quad (5)$$

respectively, corresponding to the production of $Z^+(4430)$ with $J^P = 0^-, 1^-, 2^-$. Here p_a, p_b, p_1 and p_2 denote the four-momentum of $p, \bar{p}, Z^+(4430)$ and π^- respectively.

Since $Z^+(4430)$ is observed in the invariant mass spectrum of $\psi'\pi^+$; thus, we further explore $p\bar{p} \rightarrow Z^+(4430)\pi^- \rightarrow \psi'\pi^+\pi^-$ process, depicted in Fig. 2.




FIG. 2: The diagram depicts $p\bar{p} \rightarrow \pi^- Z^+(4430) \rightarrow \pi^- \psi' \pi^+$.

The interactions of $Z^+(4430)$ with π^- and ψ' are

$$\begin{aligned} \mathcal{L}_{Z^+\pi^-\psi'} &= \begin{cases} ig_{Z\pi\psi'}^{(0)} (\pi^- \partial_\mu Z - \partial_\mu \pi^- Z) \psi'^\mu & \text{for } J^P = 0^- \\ ig_{Z\pi\psi'}^{(1)} \varepsilon_{\mu\nu\alpha\beta} \partial^\mu \psi'^\nu \partial^\alpha Z^\beta \pi^- + h.c. & \text{for } J^P = 1^- \\ ig_{Z\pi\psi'}^{(2)} \psi'_\mu Z^{\mu\nu} \partial_\nu \pi^- & \text{for } J^P = 2^- \end{cases} \end{aligned} \quad (6)$$

The coupling constants $g_{Z\pi\psi'}^{(0),(1),(2)}$ will be discussed in Sec. IV. It is straightforward to obtain the amplitudes:

$$\begin{aligned} \mathcal{M}^{J=0} &= \sqrt{2}\mathcal{G}g_{pnZ}^{(0)}g_{Z\pi^+\psi'}^{(0)}\bar{v}_{\bar{p}}\gamma_5\frac{1}{\not{p}_3 - \not{p}_b - m_n}\gamma_5u_p \\ &\times \frac{1}{(q^2 - m_Z^2 + im_Z\Gamma_Z)}(p_1 + 2p_2) \cdot \varepsilon, \end{aligned} \quad (7)$$

$$\begin{aligned} \mathcal{M}^{J=1} &= \sqrt{2}i\mathcal{G}g_{pnZ}^{(1)}g_{Z\pi^+\psi'}^{(1)}\bar{v}_{\bar{p}}\gamma_5\frac{1}{\not{p}_3 - \not{p}_b - m_n}\gamma_\nu u_p \\ &\times \frac{g_{\nu\beta} - q_\nu q_\beta/q^2}{(q^2 - m_Z^2 + im_Z\Gamma_Z)}\varepsilon_{\rho\mu\alpha\beta}p_1^\rho\varepsilon^\mu(p_1 + p_2)^\alpha, \end{aligned} \quad (8)$$

$$\begin{aligned} \mathcal{M}^{J=2} &= -\sqrt{2}\mathcal{G}g_{pnZ}^{(2)}g_{Z\pi^+\psi'}^{(2)}\bar{v}_{\bar{p}}\gamma_5\frac{1}{\not{p}_3 - \not{p}_b - m_n}\gamma_5u_p \\ &\times \frac{1}{(q^2 - m_Z^2 + im_Z\Gamma_Z)}p_a^\alpha(p_3 - p_b)^\beta\varepsilon^\mu p_2^\nu \\ &\times \left(\frac{q_{\mu\alpha}q_{\nu\beta} + q_{\mu\beta}q_{\nu\alpha}}{2} - \frac{q_{\mu\nu}q_{\alpha\beta}}{3}\right) \end{aligned} \quad (9)$$

respectively corresponding to the production of $Z^+(4430)$ with $J^P = 0^-, 1^-, 2^-$. Here $q = q_1 + q_2$ and $q_{ab} = -g_{ab} + q_a q_b/q^2$.

The cross section of $p\bar{p} \rightarrow \psi'\pi^+\pi^-$ with intermediate state $Z^+(4430)$, which involves three final states ψ', π^+

and π^- , can be obtained by integrating over the phase space [24]

$$\begin{aligned} \sigma(p\bar{p} \rightarrow \psi' \pi^+ \pi^-) \\ = \int_{a_1}^{a_2} dp_2^0 \int_{b_1}^{b_2} dp_3^0 \int_0^{2\pi} d\eta \int_{-1}^1 d(\cos \theta) d\sigma, \end{aligned} \quad (10)$$

where a_1 , a_2 , b_1 and b_2 are defined as respectively

$$\begin{aligned} a_1 &= m_2, \quad a_2 = \frac{\sqrt{s}}{2} - \frac{(m_1 + m_3)^2 - m_2^2}{2\sqrt{s}}, \\ b_1 &= \frac{1}{2\tau} \left[\rho(\tau + m_+ m_-) - |\mathbf{p}_2| \sqrt{(\tau - m_+^2)(\tau - m_-^2)} \right], \\ b_2 &= \frac{1}{2\tau} \left[\rho(\tau + m_+ m_-) + |\mathbf{p}_2| \sqrt{(\tau - m_+^2)(\tau - m_-^2)} \right], \\ \rho &= \sqrt{s} - p_2^0, \quad \tau = \rho^2 - |\mathbf{p}_2|^2, \quad m_{\pm} = m_3 \pm m_1. \end{aligned}$$

The definition of $d\sigma$ reads as

$$d\sigma = \frac{1}{32(2\pi)^4 |\mathbf{p}_a| \sqrt{s}} |\mathcal{M}|^2.$$

Here $m_{1,2,3}(p_{1,2,3})$ denote the masses (four-momentum) of ψ' , π^+ and π^- , respectively. $\mathcal{M}$ represents the amplitude shown in (7-9).

III. THE PRODUCTION OF $Z^-(4430)$ IN THE ANTIPROTON-NEUTRON ANNIHILATION PROCESS

In this section, we study the process that $\bar{p}n \rightarrow Z^-(4430) \rightarrow \psi' + \pi^-$, which is depicted in Fig. 3. Figure




FIG. 3: The production of $Z^+(4430)$ by antiproton-neutron annihilation.

4 describes the background $\bar{p}n \rightarrow \psi' \pi^-$ relevant to the production of $Z^+(4430)$.

Thus, we can directly obtain the amplitudes for the production of $Z^+(4430)$ at antiproton-neutron annihilation, which are represented as

$$\begin{aligned} \mathcal{M}^{J=0} &= g_{Zpn}^{(0)} g_{Z\pi\psi'}^{(0)} \bar{v}_{\bar{p}} \gamma_5 u_n \\ &\times \frac{1}{(q^2 - m_Z^2 + im_Z \Gamma_Z)} (p_1 + 2p_2) \cdot \varepsilon, \end{aligned} \quad (11)$$




FIG. 4: The background of the production of $Z^+(4430)$ in the antiproton-neutron interaction process.

$$\begin{aligned} \mathcal{M}^{J=1} &= ig_{Zpn}^{(1)} g_{Z\pi\psi'}^{(1)} \bar{v}_{\bar{p}} \gamma_\nu u_n \\ &\times \frac{g_{\nu\beta} - q_\nu q_\beta / q^2}{(q^2 - m_Z^2 + im_Z \Gamma_Z)} \epsilon_{\rho\mu\alpha\beta} p_1^\rho \varepsilon^\mu (p_1 + p_2)^\alpha, \end{aligned} \quad (12)$$

$$\begin{aligned} \mathcal{M}^{J=2} &= -g_{Zpn}^{(2)} g_{Z\pi\psi'}^{(2)} \bar{v}_{\bar{p}} \gamma_5 u_n \\ &\times \frac{1}{(q^2 - m_Z^2 + im_Z \Gamma_Z)} p_a^\alpha p_b^\beta \varepsilon^\mu p_2^\nu \\ &\times \left(\frac{q_{\mu\alpha} q_{\nu\beta} + q_{\mu\beta} q_{\nu\alpha}}{2} - \frac{q_{\mu\nu} q_{\alpha\beta}}{3} \right) \end{aligned} \quad (13)$$

respectively corresponding to the production of $Z^+(4430)$ with $J^P = 0^-, 1^-, 2^-$. Here $q = q_1 + q_2$ and $q_{\mu\nu} = -g_{\mu\nu} + q_\mu q_\nu / q^2$.

For obtaining the amplitude of the background $\bar{p}n \rightarrow \psi' \pi^-$ shown in Fig. 4, we need the effective Lagrangian of the strong interaction between ψ' and the nucleons

$$\mathcal{L}_{\bar{N}N\psi'} = ig_{p\bar{p}\psi'} p \gamma^\mu \bar{p} \psi'_\mu + ig_{n\bar{n}\psi'} n \gamma^\mu \bar{n} \psi'_\mu. \quad (14)$$

Since the partial decay width of $\psi' \rightarrow p\bar{p}$ is given experimentally, the coupling constant $g_{p\bar{p}\psi'}$ can be obtained. In Ref. [19], Barnes and Li obtained $g_{p\bar{p}\psi'} = (0.97 \pm 0.04) \times 10^{-3}$ by the experimental branching ratio $B(\psi' \rightarrow p\bar{p}) = (2.65 \pm 0.22) \times 10^{-4}$. Due to SU(2) symmetry, one can estimate the relation of $g_{n\bar{n}\psi'}$ and $g_{p\bar{p}\psi'}$ by $g_{n\bar{n}\psi'} \approx g_{p\bar{p}\psi'}$.

The amplitudes of the process $\bar{p}n \rightarrow \psi' \pi^-$ read as

$$\mathcal{M}_a = \sqrt{2} i \mathcal{G} g_{n\psi'} \bar{v}_{\bar{p}} \gamma_5 \frac{1}{\not{p}_a - \not{p}_2 - m_n} \not{\varepsilon} u_n, \quad (15)$$

$$\mathcal{M}_b = \sqrt{2} i \mathcal{G} g_{p\psi'} \bar{v}_{\bar{p}} \not{\varepsilon} \frac{1}{\not{p}_a - \not{p}_1 - m_p} \gamma_5 u_n. \quad (16)$$

IV. NUMERICAL RESULT AND DISCUSSION

The input parameters include $m_{\psi'} = 3686.1$ MeV, $m_{\pi^\pm} = 139.6$ MeV, $m_p = 938.3$ MeV, $m_n = 939.6$ MeV [25]; $m_Z = 4433$ MeV, $\Gamma_Z = 44$ MeV [1].

In [3], Meng and Chao calculated the decay width of $Z^+(4430) \rightarrow \psi' \pi^+$ by rescattering mechanism assuming $Z^+(4430)$ to be an S-wave $D_1 \bar{D}^*$ molecular state, and their result indicates that the order of magnitude of $\Gamma_{Z\psi'\pi^+}$ is several MeV, which can be applied to estimate the coupling constants $g_{Z\pi\psi'}^{(0,1,2)}$. Thus, in this work we

adopt the below coupling constants

$$g_{Z\pi\psi'}^{(0)} = 0.08, \quad g_{Z\pi\psi'}^{(1)} = 0.05 \text{ GeV}^{-1}, \quad g_{Z\pi\psi'}^{(2)} = 0.33,$$

which are determined by taking $\Gamma_{Z\psi'\pi^+} = 1 \text{ MeV}$.




FIG. 5: The cross section of $p\bar{p} \rightarrow Z^+(4430)\pi^-$ process with three possible quantum numbers $J^P = 0^-, 1^-, 2^-$ for $Z^+(4430)$.




FIG. 6: The cross section of $p\bar{p} \rightarrow \psi'\pi^+\pi^-$ with the intermediate state $Z^+(4430)$.

It is difficult to obtain the value of the coupling constant $g_{Z^+p\bar{n}}$ by a reliable dynamics calculation. On taking the total width of $Z^+(4430)$ as the maximum of the $\Gamma_{Z^+ \rightarrow p\bar{n}}$, one gets the maximum of coupling constant of $Z^+p\bar{n}$

$$g_{Zpn}^{(0)} = 0.52, \quad g_{Zpn}^{(1)} = 1.27, \quad g_{Zpn}^{(2)} = 0.36 \text{ GeV}^{-2}.$$

Using the above parameters, we obtain the dependence of the cross section of $p\bar{p} \rightarrow Z^+(4430)\pi^-$ on the center-of-mass energy $\sqrt{s}$, which is shown in Fig. 5.

The design luminosity of PANDA is $2 \times 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$ for the collision of the antiproton and the proton tar-

get [18]. Thus, the integrated luminosity is able to reach approximately $1.5 \text{ fb}^{-1}/\text{year}$ if the machine is of 50% efficiency and in half a year of data taking [20]. There exist sizable events for the production of $Z^+(4430)\pi^-$ in $p\bar{p}$ scattering, i.e. the maximum events with $\sqrt{s} = 5 \text{ GeV}$ can be up to 8.9×10^{10} , 1.4×10^{12} and 2.9×10^{12} , corresponding to $Z^+(4430)$ productions with $J^P = 0^-, 1^-, 2^-$, respectively. However, one needs to note that the above observation is only valid on taking the total width of $Z^+(4430)$ as the decay width of $Z^+(4430) \rightarrow p\bar{n}$. If the order of magnitude of $\Gamma_{Z^+p\bar{n}}$ is about 1 MeV, the above estimated events is suppressed by a factor $(1/44)^2 \sim$

5×10^{-4} , still enough to comply with the needs of the experimentalist. Even though $\Gamma_{Z^+ p\bar{n}}$ is at the order of keV, there also exists $10 \sim 10^2$ events of $Z^+(4430)$ production.

In Fig. 6, we present the variation of the cross section of $p\bar{p} \rightarrow Z^+(4430)\pi^- \rightarrow \psi'\pi^+\pi^-$ to $\sqrt{s}$. Due to $p\bar{p} \rightarrow \psi'\pi^+\pi^-$ with intermediate state $Z^+(4430)$ being a $2 \rightarrow 3$ process, the cross section is 3 $\sim$ 4 order smaller than that of $p\bar{p} \rightarrow Z^+(4430)\pi^-$. The line shape of the cross section of $p\bar{p} \rightarrow \pi^- Z^+(4430) \rightarrow \psi'\pi^+\pi^-$ is similar to that of $p\bar{p} \rightarrow Z^+(4430)\pi^-$.

Being a low-energy collision experiment of the antiproton with fixed target, the PANDA experiment has unique advantages to search the $Z^\pm(4430)$. Based on the above numerical analysis, we suggest the forthcoming PANDA experiment to carry out the search of $Z^\pm(4430)$, which will be helpful to finally establish $Z^\pm(4430)$ and shed light on the nature of $Z^\pm(4430)$ further.

Figure 7 shows the cross sections of $\bar{p}n \rightarrow Z^-(4430) \rightarrow$

$\psi'\pi^-$ as the function of $\sqrt{s}$, which includes the contribution from the background. Here we present the different results by taking several values of the coupling constant $g_{Zp\bar{n}}$, which correspond to the $\Gamma_{Z\bar{p}n} = 44, 22, 4.4$ and 1 MeV respectively. The line shape of the cross section of $\bar{p}n \rightarrow Z^-(4430) \rightarrow \psi'\pi^-$ shows the enhancement at energy $\sqrt{s} = 4.433$ GeV, which is different from the case of $Z^+(4430)$ production at $p\bar{p}$ scattering process.

In terms of our calculation, we find that the largest production rate of $Z^+(4430)$ mainly comes from the $p\bar{n}$ annihilation process. Unfortunately, the PANDA experiment mainly refers to the collision of proton and antiproton. Thus, it is difficult to test our proposal of $Z^+(4430)$ production by $p\bar{n}$ annihilation based on present high-energy experiments. One of the difficulties is that it is not easy to provide and control the source of the neutron, which is waiting to be solved by clever experimentalists.




FIG. 7: The cross section of the production of $Z^-(4430)$ in $\bar{p}n$ annihilation process.

Acknowledgments

One of us (X.L.) would like to thank Prof. Eef van Beveren for useful discussion and fruitful comments. This work was supported by the *Fundação para a Ciência e*

a Tecnologia of the Ministério da Ciência, Tecnologia e Ensino Superior of Portugal (SFRH/BPD/34819/2007) and the National Natural Science Foundation of China under Grant 10705001.

-
- [1] Belle Collaboration, K. Abe et al., Phys. Rev. Lett. **100**, 142001 (2008).
 - [2] J.L. Rosner, Phys. Rev. **D 76**, 114002 (2007).
 - [3] C. Meng and K.T. Chao, arXiv:0708.4222 [hep-ph].
 - [4] X. Liu, Y.R. Liu, W.Z. Deng and S.L. Zhu, Phys. Rev. **D 77**, 034003 (2008), arXiv:0711.0494.
 - [5] X. Liu, Y.R. Liu, W.Z. Deng, S.L. Zhu, Phys. Rev. **D 77**, 094015 (2008), arXiv:0803.1295 [hep-ph].
 - [6] G.J. Ding, arXiv:0711.1485 [hep-ph]; G.J. Ding, W. Huang, J.F. Liu and M.L. Yan, arXiv:0805.3822 [hep-ph].
 - [7] S.H. Lee, A. Mihara, F.S. Navarra and M. Nielsen, arXiv:0710.1029 [hep-ph].
 - [8] L. Maiani, A.D. Polosa and V. Riquer, arXiv:0708.3997 [hep-ph].
 - [9] K. Cheung, W.Y. Keung and T.C. Yuan, Phys. Rev. **D 76**, 117501 (2007).
 - [10] S.S. Gershtein, A.K. Likhoded and G.P. Pronko, arXiv:0709.2058 [hep-ph].
 - [11] D.V. Bugg, arXiv:0802.0934 [hep-ph].
 - [12] C.F. Qiao, arXiv:0709.4066 [hep-ph].
 - [13] E. Braaten and M. Lu, arXiv:0712.3885 [hep-ph].
 - [14] Y. Li, C.D. Lu and W. Wang, Phys. Rev. **D 77**, 054001 (2008).
 - [15] X.H. Liu, Q. Zhao and F.E. Close, arXiv:0802.2648 [hep-ph].
 - [16] X. Liu, B. Zhang and S.L. Zhu, Phys. Rev. **D 77**, 114021, (2008), arXiv:0803.4270 [hep-ph].
 - [17] T. Matsuki, T. Morii and K. Sudoh, arXiv:0805.2442 [hep-ph].
 - [18] PANDA Collaboration, Technical Progress Report for: PANDA, <http://www-panda.gsi.de>.
 - [19] T. Barnes and X. Li, Phys. Rev. **D 75**, 054018 (2007).
 - [20] G.Y. Chen and J.P. Ma, arXiv:0802.2982 [hep-ph].
 - [21] M.K. Gaillard, L. Maiani and R. Petronzio, Phys. Lett. **B 110**, 489 (1982).
 - [22] A. Lundborg, T. Barnes and U. Wiedner, Phys. Rev. **D 73**, 096003 (2006).
 - [23] Z. Lin, C.M. Ko and B. Zhang, Phys. Rev. **C 61**, 024904 (2000); B. Holzenkamp et al. Nucl.Phys. **A 500**, 485 (1989).
 - [24] T. Hahm, *FormCalc 5 Users Guide*, <http://www.feynarts.de/formcalc>.
 - [25] W.M. Yao et al., Particle Data Group, J. Phys. G **33**, 1 (2006).